


Introduction


- NMR is the most powerful tool available for organic structure determination.
- It is used to study a wide variety of nuclei:
 - ^1H
 - ^{13}C
 - ^{15}N
 - ^{19}F
 - ^{31}P

=>




Nuclear Spin


- A nucleus with an odd atomic number or an odd mass number has a nuclear spin.
- The spinning charged nucleus generates a magnetic field.



spinning proton




loop of current




bar magnet

=>




External Magnetic Field

When placed in an external field, spinning protons act like bar magnets.



=>




Two Energy States


The magnetic fields of the spinning nuclei will align either *with* the external field, or *against* the field.

A photon with the right amount of energy can be absorbed and cause the spinning proton to flip.


=>



ΔE and Magnet Strength



- Energy difference is proportional to the magnetic field strength.
- $$\Delta E = h\nu = \frac{\gamma \hbar B_0}{2\pi}$$
- Gyromagnetic ratio, γ , is a constant for each nucleus ($26,753 \text{ s}^{-1}\text{gauss}^{-1}$ for H).
- In a 14,092 gauss field, a 60 MHz photon is required to flip a proton.
- Low energy, radio frequency. \Rightarrow




Magnetic Shielding


- If all protons absorbed the same amount of energy in a given magnetic field, not much information could be obtained.
- But protons are surrounded by electrons that shield them from the external field.
- Circulating electrons create an induced magnetic field that opposes the external magnetic field.


=>

Shielded Protons




Magnetic field strength must be increased for a shielded proton to flip at the same frequency.






Protons in a Molecule

Depending on their chemical environment, protons in a molecule are shielded by different amounts.




=>




NMR Signals



- The ***number*** of signals shows how many different kinds of protons are present.
- The ***location*** of the signals shows how shielded or deshielded the proton is.
- The ***intensity*** of the signal shows the number of protons of that type.
- Signal ***splitting*** shows the number of protons on adjacent atoms. =>




The NMR Spectrometer




The NMR Graph






Tetramethylsilane



- TMS is added to the sample.
- Since silicon is less electronegative than carbon, TMS protons are highly shielded. Signal defined as zero.
- Organic protons absorb downfield (to the left) of the TMS signal.

=>




Chemical Shift



- Measured in parts per million.
- Ratio of shift downfield from TMS (Hz) to total spectrometer frequency (Hz).
- Same value for 60, 100, or 300 MHz machine.
- Called the delta scale.


=>

Delta Scale



$$\text{chemical shift, ppm } \delta = \frac{\text{shift downfield from TMS (in Hz)}}{\text{spectrometer frequency (in MHz)}}$$





Location of Signals

TABLE 13-2 Chemical Shifts of the Chloromethanes


Compound	Chemical Shift	Difference
$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array}$	$\delta 0.2$	
$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{Cl} \\ \\ \text{H} \end{array}$	$\delta 3.0$	2.8 ppm
$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{Cl} \\ \\ \text{Cl} \end{array}$	$\delta 5.3$	2.3 ppm
$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{Cl} \\ \\ \text{Cl} \\ \\ \text{Cl} \end{array}$	$\delta 7.2$	1.9 ppm

Note: Each chlorine atom added changes the chemical shift of the remaining methyl protons by about 2 to 3 ppm. These changes are nearly additive.

- More electronegative atoms deshield more and give larger shift values.
- Effect decreases with distance.
- Additional electronegative atoms cause increase in chemical shift.

=>



Typical Values



Type of Proton	Approximate δ	Type of Proton	Approximate δ
alkane ($-\text{CH}_3$)	0.9	>C=C<CH_3	1.7
alkane ($-\text{CH}_2-$)	1.3	Ph—H	7.2
alkane ($-\text{CH}-$)	1.4	Ph—CH ₃	2.3
$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{CH}_3 \end{array}$	2.1	R—CHO	9–10
$-\text{C}\equiv\text{C}-\text{H}$	2.5	R—COOH	10–12
R—CH ₂ —X (X = halogen, O)	3–4	R—OH	variable, about 2–5
$\begin{array}{c} \text{>C=C<} \\ \text{H} \end{array}$	5–6	Ar—OH	variable, about 4–7
		R—NH ₂	variable, about 1.5–4

Note: These values are approximate, as all chemical shifts are affected by neighboring substituents. The numbers given here assume that alkyl groups are the only other substituents present. A more complete table of chemical shifts appears in Appendix 1.



Aromatic Protons, $\delta 7\text{-}\delta 8$



induced field
reinforces the
external field
(deshielding)

=>

Vinyl Protons, $\delta 5\text{-}\delta 6$





induced field reinforces the external field (deshielding)






Acetylenic Protons, $\delta 2.5$

B_{induced} shields
the proton



Aldehyde Proton, $\delta 9$ - $\delta 10$







O-H and N-H Signals

- Chemical shift depends on concentration.
- Hydrogen bonding in concentrated solutions deshield the protons, so signal is around $\delta 3.5$ for N-H and $\delta 4.5$ for O-H.
- Proton exchanges between the molecules broaden the peak.



=>

Carboxylic Acid Proton, $\delta 10+$




Number of Signals


Equivalent hydrogens have the same chemical shift.




Intensity of Signals




- The area under each peak is proportional to the number of protons.
- Shown by integral trace.




How Many Hydrogens?



When the molecular formula is known, each integral rise can be assigned to a particular number of hydrogens.







Spin-Spin Splitting



- Nonequivalent protons on adjacent carbons have magnetic fields that may align with or oppose the external field.
- This magnetic coupling causes the proton to absorb slightly downfield when the external field is reinforced and slightly upfield when the external field is opposed.
- All possibilities exist, so signal is split. =>

1,1,2-Tribromoethane

Nonequivalent protons on adjacent carbons.




Triplet: 2 Adjacent Protons



=>

The $N + 1$ Rule




If a signal is split by N equivalent protons, it is split into $N + 1$ peaks.

Relative Peak Intensities of Symmetric Multiplets		
<i>Number of Equivalent Protons Causing Splitting</i>	<i>Number of Peaks (multiplicity)</i>	<i>Area Ratios (Pascal's triangle)</i>
0	1 (singlet)	1
1	2 (doublet)	1 1
2	3 (triplet)	1 2 1
3	4 (quartet)	1 3 3 1
4	5 (quintet)	1 4 6 4 1
5	6 (sextet)	1 5 10 10 5 1
6	7 (septet)	1 6 15 20 15 6 1





Range of Magnetic Coupling



- Equivalent protons do not split each other.
- Protons bonded to the same carbon will split each other only if they are not equivalent.
- Protons on adjacent carbons normally will couple.
- Protons separated by four or more bonds will not couple.



=>

Splitting for Ethyl Groups




=>

Splitting for Isopropyl Groups



=>




Coupling Constants

- Distance between the peaks of multiplet
- Measured in Hz
- Not dependent on strength of the external field
- Multiplets with the same coupling constants may come from adjacent groups of protons that split each other.

=>


Values for Coupling Constants




		<u>Approx. J</u>		<u>Approx. J</u>
$\begin{array}{c} & \\ -C & -C- \\ & \\ H & H \end{array}$	(free rotation)	7 Hz ^a		8 Hz
			(ortho)	
$\begin{array}{c} \diagdown & \diagup \\ C & =C \\ \diagup & \diagdown \\ H & H \end{array}$	(cis)	10 Hz		2 Hz
			(meta)	
$\begin{array}{c} \diagdown & \diagup \\ C & =C \\ \diagup & \diagdown \\ H & H \end{array}$	(trans)	15 Hz		6 Hz
			(para)	
$\begin{array}{c} \diagdown & \diagup \\ C & =C \\ \diagup & \diagdown \\ & H \\ & H \end{array}$	(geminal)	2 Hz		
			(allylic)	

^aThe value of 7 Hz in an alkyl group is averaged for rapid rotation about the carbon-carbon bond. If rotation is hindered by a ring or bulky groups, other splitting constants may be observed.

⇒







Complex Splitting





- Signals may be split by adjacent protons, different from each other, with different coupling constants.
- Example: H^a of styrene which is split by an adjacent H *trans* to it ($J = 17$ Hz) and an adjacent H *cis* to it ($J = 11$ Hz).

=>

Splitting Tree




Spectrum for Styrene



=>





Stereochemical Nonequivalence




- Usually, two protons on the same C are equivalent and do not split each other.
- If the replacement of each of the protons of a $-\text{CH}_2$ group with an imaginary “Z” gives stereoisomers, then the protons are non-equivalent and will split each other.

=>

Some Nonequivalent Protons




Time Dependence




- Molecules are tumbling relative to the magnetic field, so NMR is an averaged spectrum of all the orientations.
- Axial and equatorial protons on cyclohexane interconvert so rapidly that they give a single signal.
- Proton transfers for OH and NH may occur so quickly that the proton is not split by adjacent protons in the molecule.

=>


Hydroxyl proton



- Ultrapure samples of ethanol show splitting.
- Ethanol with a small amount of acidic or basic impurities will not show splitting.




=>




N-H Proton

- Moderate rate of exchange.
- Peak may be broad.




=>

Identifying the O-H or N-H Peak



- Chemical shift will depend on concentration and solvent.
- To verify that a particular peak is due to O-H or N-H, shake the sample with D_2O
- Deuterium will exchange with the O-H or N-H protons.
- On a second NMR spectrum the peak will be absent, or much less intense.

=>




Carbon-13

- ^{12}C has no magnetic spin.
- ^{13}C has a magnetic spin, but is only 1% of the carbon in a sample.
- The gyromagnetic ratio of ^{13}C is one-fourth of that of ^1H .
- Signals are weak, getting lost in noise.
- Hundreds of spectra are taken, averaged.

=>


Fourier Transform NMR



- Nuclei in a magnetic field are given a radio-frequency pulse close to their resonance frequency.
 - The nuclei absorb energy and precess (spin) like little tops.
 - A complex signal is produced, then decays as the nuclei lose energy.
 - Free induction decay is converted to spectrum.
- =>













Hydrogen and Carbon

Chemical Shifts





-COOH


$\delta 11-12$



Combined ^{13}C and ^1H Spectra




Differences in ^{13}C Technique



- Resonance frequency is ~ one-fourth, 15.1 MHz instead of 60 MHz.
- Peak areas are not proportional to number of carbons.
- Carbon atoms with more hydrogens absorb more strongly.


=>



Spin-Spin Splitting

- It is unlikely that a ^{13}C would be adjacent to another ^{13}C , so splitting by carbon is negligible.
- ^{13}C will magnetically couple with attached protons and adjacent protons.
- These complex splitting patterns are difficult to interpret.


=>



Proton Spin Decoupling

- To simplify the spectrum, protons are continuously irradiated with “noise,” so they are rapidly flipping.
- The carbon nuclei see an average of all the possible proton spin states.
- Thus, each different kind of carbon gives a single, unsplit peak.

=>




Off-Resonance Decoupling


- ^{13}C nuclei are split only by the protons attached directly to them.
- The $N + 1$ rule applies: a carbon with N number of protons gives a signal with $N + 1$ peaks.

=>


Interpreting ^{13}C NMR




- The number of different signals indicates the number of different kinds of carbon.
- The location (chemical shift) indicates the type of functional group.
- The peak area indicates the numbers of carbons (if integrated).
- The splitting pattern of off-resonance decoupled spectrum indicates the number of protons attached to the carbon. =>



Two ^{13}C NMR Spectra






MRI

- Magnetic resonance imaging, noninvasive
- “Nuclear” is omitted because of public’s fear that it would be radioactive.
- Only protons in one plane can be in resonance at one time.
- Computer puts together “slices” to get 3D.
- Tumors readily detected.

=>



End of Chapter 13